
Citation:

Astita, R and Tashani, O and Hind, K and Sharp, D and Johnson, MI (2015) A PRELIMINARY INVESTIGATION OF THE RELATIONSHIP BETWEEN PAIN SENSITIVITY, BODY FAT DISTRIBUTION AND BLOOD LEVELS OF IL-6, CRP, TNF- AND LEPTIN. In: 9th Congress of the European Pain Federation EFIC, 02 Sep 2015 - 05 Sep 2015, Vienna, Austria.

Link to Leeds Beckett Repository record:

<https://eprints.leedsbeckett.ac.uk/id/eprint/1395/>

Document Version:

Conference or Workshop Item (Accepted Version)

The aim of the Leeds Beckett Repository is to provide open access to our research, as required by funder policies and permitted by publishers and copyright law.

The Leeds Beckett repository holds a wide range of publications, each of which has been checked for copyright and the relevant embargo period has been applied by the Research Services team.

We operate on a standard take-down policy. If you are the author or publisher of an output and you would like it removed from the repository, please [contact us](#) and we will investigate on a case-by-case basis.

Each thesis in the repository has been cleared where necessary by the author for third party copyright. If you would like a thesis to be removed from the repository or believe there is an issue with copyright, please contact us on openaccess@leedsbeckett.ac.uk and we will investigate on a case-by-case basis.

A preliminary Investigation of the relationship between pain sensitivity, body fat distribution and blood levels of IL-6, CRP, TNF- α and leptin.

Rehab Astita, Osama A. Tashani, Karen Hind, Duncan Sharp and Mark I. Johnson

Centre for Pain Research and Carnegie Research Institute, Leeds Beckett University

Abstract

Background and aims: Obese individuals have increased concentrations of pro-inflammatory cytokines and some other markers of inflammation and an increased risk of metabolic disorders. The relationship between obesity, pro-inflammatory cytokines and pain sensitivity response is not fully understood. **Aim of the Study:** To investigate associations between body fat distribution, C reactive protein (CRP), interleukin 6 (IL-6), tumour necrosis factor alpha (TNF- α), leptin and pain sensitivity in adults.

Methods: 38 adults (n=18 women) were grouped as normal weight (n=22) or obese (n=16), based on body mass index (BMI). Measurements of pressure pain and cold pressor pain sensitivity response, biomarkers (venous blood), and body composition (dual X-ray absorptiometry) were evaluated for each participant.

Results: Pressure pain threshold was significantly lower in obese (mean \pm SD=340.93 \pm 93.58 kpa) compared to the normal weight group (447.45 \pm 203.72 kpa, p=0.039, t-test). Forward regression suggested that high gynoid fat (g) was associated with lower pressure pain thresholds (β =-0.383, p=0.028), high lower limb fat (g) was associated with lower cold pain thresholds (β =-0.495, p=0.003) and high IL-6 predicted higher cold pain tolerance (β =0.345, p=0.049). Women were more sensitive to pressure pain (P=0.03).

Conclusion: Gynoid and lower limb fat content correlated with pain sensitivity response in adults, whereby those with greater fat content were more sensitive to pain. However, this may be a reflection of the sex differences in pain sensitivity as women have greater gynoid and leg fat contents.